

**Eu-Tec
Nutrition**

In Your Corner

Transition Feed



Product: Later transition diet for pigs weighing 15—25 lbs.

Pack Size: 50 lb Bags. Also available in 1,200 to 2,000 lb totes.

Form: 2 mm Pellet

Use Rate: Feed 3 lbs to 10 lbs to pigs over 15 lbs up to 25 lbs bodyweight.

Features:

- Superb 3rd stage diets for pigs weaned over 18 days of age.
- Ideal for larger pigs, but still includes rolled oats and the highest quality fish meals.
- Formulated to exact nutritional specifications
- Uses food grade ingredients to help the piglets digestive system
- 2 mm pellets are made using European technology that assures high pellet durability yet remains soft enough to be easily chewed by the young weaned pig

Purpose: Ideal 3rd stage starter diet for pigs weaned at over 12 lbs and 18 days/

Benefits:

- Highly digestible diets promote feed intake
- Improved weight gain and feed efficiency
- Fewer digestive upsets due to poor intakes and low digestibility

Remarks: Keep feed fresh. Do not store in nursery rooms.

Feeding Directions:

- Even older, weaned pigs need at least 1 lb of either sureSTART, easiStart and 3 lbs of multiStart or Thrust depending on weaning age and bodyweight.
- We recommend small quantities of fresh food frequently, not allowing feed to get stale. Any change in diet should be made over 3-4 days to minimize the risk of digestive upsets. Also, changes of food should not be within 3 days of weaning.
- It is important to give piglets an abundant, clean source of fresh water



**Eu-Tec
Nutrition**

In Your Corner

305 Pershing Rd. Maquoketa, IA 52060 Ph: 563-652-2977 www.eu-tecnutrition.com



VITAL +DG

MEDICATED

A Complete Starter Feed for Pigs

For control of swine dysentery associated with Brachyspira hyodysenteriae susceptible to Tiamulin. For treatment of swine bacteria enteritis caused by Escherichia coli and Salmonella choleraesuis and bacterial pneumonia caused by Pasteurella multocida susceptible to Chlortetracycline.

ACTIVE DRUG INGREDIENT

Tiamulin Hydrogen Fumarate	0.0039%	35g/ton
Chlortetracycline	0.044%	400 g/ton

WARNING: Withdraw medicated feed two days before slaughter.

GUARANTEED ANALYSIS

Crude Protein	19.00%	min
Lysine	1.50%	min
Crude Fat	6.50%	min
Crude Fiber	2.50%	max
Calcium (Ca)	0.50%	min
Calcium (Ca)	1.00%	max
Salt	0.75%	min
Salt	1.25%	max
Phosphorus (P)	0.60%	min
Selenium (Se)	0.3 ppm	min
Zinc (Zn)	3,000 ppm	min
Vitamin A	5,000 IU/lb	min
Vitamin D	900 IU/lb	min
Vitamin E	75 IU/lb	min

INGREDIENT STATEMENT

Corn, Soybean Meal, Ground Heat Processed Soybeans, Dried Whey Permeate, Cereal Food Fines, Steam Rolled Oat Groats, Fish Meal, Soybean Oil, Sodium Bentonite, Calcium Carbonate, Zinc Oxide, L-Lysine, Encapsulated Salt of Alimentary Fatty Acid, Encapsulated Essential Fatty Acid Mixture, Hydrated Silicone Dioxide, Salt, Mono-Calcium Phosphate, dl-Methionine, l-Threonine, Dried Bacillus Licheniformis Fermentatin Product, Dried Bacillus Subtilis Fermentation Product, dl-Tryptophan, Vitamin A Acetate, Vitamin D₃ Supplement, Vitamin E Supplement, Menadione Dimethylpyrimidinol Bisulfite (source of Vitamin K), Thiamine, Riboflavin, Pyridoxine Hydrochloride, Vitamin B₁₂ Supplement, Biotin, Calcium Pantothenate, Niacin Supplement, Folic Acid, Choline Chloride, Ethylenediamine Dihydriodide, Sodium Selenite, Selenium Yeast, Cobalt Carbonate, Iron Sulfate, Iron Protein-ate, Manganese Sulfate, Copper Sulfate, Zinc Sulfate, Mineral Oil.

FEEDING DIRECTIONS

Vital is the ideal diet for piglets weighing over 15 lbs. Feed continuously as the sole ration for 14 days to provide chlortetracycline at 10 mg/lb of body weight. Feed to pigs weighing from 12 to 50 pounds. Provide fresh clean water at all times.

CAUTION: Follow label directions. Do not feed to animals other than swine. Not for use in in swine weighing over 250 lbs. Use as the only source of Tiamulin. Swine being treated with Tiamulin should not have access to feeds containing polyether ionophores (e.g., Monensin, Lasalocid, Narasin, Salinomycin and Semduramicin) as adverse reactions may occur. If signs of toxicity occur, discontinue use of medicated feed.

SPECIAL CONSIDERATIONS: Protect medicated feeds from excessive temperatures. Do not store at warm temperatures, 104°F (40°C), for more than 7 days.

- Manufactured By-
Eu-Tec Nutrition, Inc.
305 Pershing Road
Maquoketa, IA 52060

NET WT 50 LB (22.68 Kg)

